

# New England Fertility Institute

1275 Summer Street • Suite 201 • Stamford, CT 06905 • Tel: (203)325-3200 • Fax: (203)323-3130  
 9 Washington Avenue • Hamden, CT 06518 • Tel: (203) 248-2353 • Fax: (203) 248-2074  
 4 Corporate Drive • Suite 286 • Shelton, CT 06484 • Tel: (203) 929-6412 • Fax: (203) 929-6428

## SUNDAY RETRIEVAL

	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY	SUNDAY
<b>GANIRELIX</b>	Ganirelix	Ganirelix	Ganirelix	<del>                    </del>	<del>                    </del>
<b>LUPRON</b>			<del>                    </del>	<del>                    </del>	<del>                    </del>
<b>LO-DOSE HCG</b>			<del>                    </del>	<del>                    </del>	<del>                    </del>
<b>MENOPUR</b>			<del>                    </del>	<del>                    </del>	<del>                    </del>
<b>FSH</b> Follistim/Gonal-F			<del>                    </del>	<del>                    </del>	HCG URINE TEST
<b>HCG</b> Chorionic Gonadotropin/ Novarel/Pregnyl	<del>                    </del>	<del>                    </del>	HCG 5,000 UNITS OR 10,000 UNITS. WE WILL CALL WITH TIME. SET YOUR ALARM.	<del>                    </del>	NOTHING TO EAT OR DRINK IN MORNING
<b>ESTRACE</b>	<u>Estrace</u> 0.5 mg 1 tab twice a day	CONTINUE	CONTINUE	CONTINUE	CONTINUE
<b>MEDROL</b>	<del>                    </del>	<del>                    </del>	<del>                    </del>	<u>Medrol</u> 16 mg 1 tab a day for 5 days	CONTINUE
<b>DOXYCYCLINE</b>	<del>                    </del>	<del>                    </del>	<del>                    </del>	<u>Doxycycline</u> 100 mg 1 tab 2 times a day for 5 days	CONTINUE
			NO SEXUAL RELATIONS	NOTHING TO EAT OR DRINK AFTER MIDNIGHT	<b>RETRIEVAL</b> ARRIVE 45 MINS EARLY WITH PARTNER

**HCG 5,000 UNITS:** Put two cc(ml) water into full powder - draw out one cc(ml) diluted medication - store remaining one cc(ml) in refrigerator.

**HCG 10,000 UNITS:** Put one cc(ml) water into full powder - draw out all the diluted powder in the one cc. Inject mixed HCG at the instructed time.

Mix with 22 gauge 1 1/2" (black) or 18 gauge 1 1/2" (pink). **Switch** to 27 gauge 1/2" (grey) or 25 gauge 5/8" (blue) to inject in belly or front of thighs.