

New England Fertility Institute

1275 Summer Street • Suite 201 • Stamford, CT 06905 • Tel: (203)325-3200 • Fax: (203)323-3130
 9 Washington Avenue • Hamden, CT 06518 • Tel: (203) 248-2353 • Fax: (203) 248-2074
 4 Corporate Drive • Suite 286 • Shelton, CT 06484 • Tel: (203) 929-6412 • Fax: (203) 929-6428

THURSDAY RETRIEVAL

	SUNDAY	MONDAY	TUESDAY	WEDNESDAY	THURSDAY
GANIRELIX	Ganirelix	Ganirelix	Ganirelix	 	
LUPRON	 	 	 	 	
LO-DOSE HCG	 	 	 	 	
MENOPUR	 	 	 	 	
FSH Follistim/Gonal-F	 	 	 	 	HCG URINE TEST
HCG Chorionic Gonadotropin/ Novarel/ Pregnyl	 	 	HCG 5,000 UNITS OR 10,000 UNITS. WE WILL CALL WITH TIME. SET YOUR ALARM.	 	NOTHING TO EAT OR DRINK IN MORNING
ESTRACE	<u>Estrace</u> 0.5 mg 1 tab twice a day	CONTINUE	CONTINUE	CONTINUE	CONTINUE
MEDROL	 	 	 	<u>Medrol</u> 16 mg 1 tab a day for 5 days	CONTINUE
DOXYCYCLINE	 	 	 	<u>Doxycycline</u> 100 mg 1 tab 2 times a day for 5 days	CONTINUE
			NO SEXUAL RELATIONS	NOTHING TO EAT OR DRINK AFTER MIDNIGHT	RETRIEVAL ARRIVE 45 MINS EARLY WITH PARTNER

HCG 5,000 UNITS: Put two cc(ml) water into full powder - draw out one cc(ml) diluted medication - store remaining one cc(ml) in refrigerator.

HCG 10,000 UNITS: Put one cc(ml) water into full powder - draw out all the diluted powder in the one cc. Inject mixed HCG at the instructed time.

Mix with 22 gauge 1 1/2" (black) or 18 gauge 1 1/2" (pink). **Switch** to 27 gauge 1/2" (grey) or 25 gauge 5/8" (blue) to inject in belly or front of thighs.